



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I. SOLUTION BY WILLIAM HOOVER, Columbus, Ohio.

Take ABC as the triangle of reference, and the trilinear coördinates of the vertices A, B, C to be $(\alpha_2, 0, 0)$; $(0, \beta_2, 0)$; $(0, 0, \gamma_2)$, and those of O , $(\alpha_1, \beta_1, \gamma_1)$; then the equation of AO is $\gamma_1\beta - \beta_1\gamma = 0$.


The coördinates of A' will be proportional to $(0, \beta_1, \gamma_1)$, and, by symmetry, those of B' to $(\alpha_1, 0, \gamma_1)$, and of C' to $(\alpha_1, \beta_1, 0)$.

The equation of $A'B'$ is $\beta_1\gamma_1\alpha + \alpha_1\gamma_1\beta - \alpha_1\beta_1\gamma = 0$.

The point of intersection of AB and $A'B'$, or Z , has coördinates proportional to $(-\alpha_1, \beta_1, 0)$; those of the intersection BC and $B'C'$, or X , to $(0, -\beta_1, \gamma_1)$; and of CA and $C'A'$, or Y , to $(\alpha_1, 0, -\gamma_1)$; and X, Y, Z are collinear, since

$$\begin{vmatrix} 0, & -\beta_1, & \gamma_1 \\ \alpha_1, & 0, & -\gamma_1 \\ -\alpha_1, & \beta_1, & 0 \end{vmatrix} = \alpha_1\beta_1\gamma_1 \begin{vmatrix} 0, & -1, & 1 \\ 1, & 0, & -1 \\ -1, & 1, & 0 \end{vmatrix}$$

$$= \alpha_1\beta_1\gamma_1 \begin{vmatrix} 0, & 0, & 1 \\ 1, & -1, & -1 \\ -1, & 1, & 0 \end{vmatrix} = \alpha_1\beta_1\gamma_1 \begin{vmatrix} 1, & -1 \\ -1, & 1 \end{vmatrix} = 0.$$



The equation of AX is $\gamma_1\beta + \beta_1\gamma = 0$, and those of BY and CZ respectively $\gamma_1\alpha + \alpha_1\gamma = 0$, and $\beta_1\alpha + \alpha_1\beta = 0$.

The coördinates of C'' , the point of intersection of AX and BY , are proportional to $(\alpha_1, \beta_1, \gamma_1)$; of BY and CZ , to $(\alpha_1, \beta_1, \gamma_1)$; and of CZ and AX to $(\alpha_1, \beta_1, \gamma_1)$, the latter two points being A'' , B'' respectively.

It is evident now that the equations to AA'' , BB'' , CC'' are the same as those of AO , BO , CO , in order, the first three lines then passing through O .

II. SOLUTION BY H. L. OLSON, Chicago, Illinois.

I shall amplify this theorem by proving that the lines AA'' , BB'' , CC'' are identical with the lines AA' , BB' , CC' respectively, and hence intersect in the point O . In the triangles $BC'Y$ and $CB'Z$, the lines BC , $C'B'$, and YZ , joining corresponding vertices, meet at X , and hence the points A, A', A'' , in which corresponding sides meet, are collinear. Similarly B, B', B'' are collinear; also C, C', C'' . Hence the lines AA'' , BB'' , CC'' meet at O .

Also solved by A. PELLETIER and the Proposer.

2716 [June, 1918]. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.

To a passenger in a train moving at the rate of 40 miles per hour, the rain appears to be rushing downward and towards him at an angle of 20 degrees with the horizontal. If the rain is actually falling in a vertical direction, show that the velocity of the raindrops in feet per second is 21.35.

SOLUTION BY E. H. WORTHINGTON, Elkins Park, Pa.

A velocity of 40 miles per hour is the same as $58\frac{2}{3}$ feet per second. If v is the velocity of the raindrop, we have $v = 58\frac{2}{3} \tan 20^\circ = 58\frac{2}{3} \times 0.364$ feet per sec. = 21.35 feet per second.

Also solved by H. E. CARLETON, A. M. HARDING, H. L. OLSON, A. PELLETIER and J. B. REYNOLDS.

2735 [December, 1918]. Proposed by H. B. PHILLIPS, Massachusetts Institute of Technology.

If two lines AE and BD , drawn from the vertices A and B of a triangle to the opposite sides, divide the angles A and B so that the parts of A are respectively less than the corresponding parts of B , then AE is greater than BD .